

Crystal structure of 8,10-diiodo-3-(iodomethyl)-2,3-dihydro-[1,4]oxazino-[2,3,4-*ij*]quinolin-4-ium triiodide, $[\text{C}_{12}\text{H}_9\text{I}_3\text{NO}] \cdot \text{I}_3$, $\text{C}_{12}\text{H}_9\text{I}_6\text{NO}$


Vladimir I. Batalov^I, Alla Dikhtiarenko^{II}, Irina D. Yushina^I, Ekaterina V. Bartashevich^I, Dimitri G. Kim^I and Santiago García-Granda^{*,III}

^I Department of Organic Chemistry, South Ural State University, 76, Lenin avenue, Chelyabinsk, Russia, 454080

^{II} Departamento Química Orgánica e Inorgánica, Universidad de Oviedo, 33006 Oviedo, Spain

^{III} Departamento de Química Física y Analítica, Facultad de Química, Universidad de Oviedo - CINN, C/ Julián Clavería, 8, 33006 Oviedo, (Asturias) Spain

Received January 20, 2014, accepted May 14, 2014, available online May 28, 2014, CCDC no. 1005550



Abstract

$\text{C}_{12}\text{H}_9\text{I}_6\text{NO}$, monoclinic, $P2_1/n$ (no. 14), $a = 9.2855(2)$ Å, $b = 17.1834(4)$ Å, $c = 12.2416(3)$ Å, $\beta = 93.258(2)^\circ$, $V = 1950.1$ Å³, $Z = 4$, $R_{\text{gt}}(F) = 0.0433$, $wR_{\text{ref}}(F^2) = 0.0630$, $T = 120$ K.

Table 1. Data collection and handling.

Crystal:	brown rhombohedrals, size $0.0131 \times 0.0460 \times 0.0846$ mm
Wavelength:	Mo K_α radiation (0.71073 Å)
μ :	95.53 cm ⁻¹
Diffractometer, scan mode:	Xcalibur, Ruby, Gemini, ω
$2\theta_{\text{max}}$:	62.74°
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$:	17314, 5893
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2\sigma(I_{\text{obs}})$, 4561
$N(\text{param})_{\text{refined}}$:	181
Programs:	CrysAlis PRO [10], SIR92 [11], SHELX [12], DIAMOND [13], WinGX [14], enCIFer [15]

Source of material

5,7-diiodo-8-(allyloxy)quinoline was obtained using 5,7-diiodo-8-quinolinol and allyl bromide in acetone/ K_2CO_3 with reflux. The solution of 5,7-diiodo-8-(allyloxy)quinoline (0.110 g, 0.25 mmol) in 10 ml of acetone was mixed with a solution of iodine (0.192 g, 0.75 mmol) in 10 ml of acetone. Single crystals for the X-ray diffraction study were obtained after keeping the resulting mixture at room temperature in closed flask for 5 days and then by slow solvent evaporation, yield: 0.135 g (57%).

Experimental details

The positions of the H atoms were calculated based on geometric criteria (C–H = 0.97 Å, 0.93 Å and 0.85 Å for methylene, aromatic and methine atoms, respectively) than have been placed in their calculated position and refined isotropically using a rider model with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Discussion

A number of synthesis for quinoline compounds with fused N-1/C-8 centers [1–3] and X-ray structures of condensed tricyclic azines [4–5] were reported. Represented crystal structure can be an interesting object for studying the directed self-assembly of polyiodide chains [6–8]. The asymmetric unit of the title compound consists of one iodoquinolinium derivative cation and one triiodide anion. The atoms N1 and C1–C9, defining the quinolinium part of molecular cation, are situated in the same plane. The C–C bonds lengths range from 1.363(7) to 1.423(7) Å; the C–N bond distances range from 1.339(7) to 1.382(6) Å. The I1 and I2 atoms are located in the same plane as the quinolinium moiety with C5–I1 and C3–I2 bond distances of 2.090(5) and 2.091(5) Å, respectively. The crystal structure confirms the closure of the six-member ring between O1 and N1, during cyclization reaction, resulted in C11–N1 bond formation. The C11–N1 bond has a covalent character (1.493(6) Å). For six-membered folded ring the C–O bond distances range from 1.351(6) to 1.432(6) Å, and C10–C11 bond length is 1.507(7) Å. In this ring the C10 atom of methylene group is significantly out of plane (24.2°). The exocyclic C11–C12 bond has a single bond character at a bond distance of 1.525(7) Å. The C12–I3 bond length is 2.146(5) Å and significantly deviated from mean plane of iodoquinolinium moiety (30.9°); the C11–C12–I3 angle is 110.5(3)°. The molecular conformation is established by intramolecular contacts C12–H12B...O1 and C10–H10A...I3 with D...A distances 2.955(6) and 3.524(5) Å, respectively. The triiodide anion is almost linear with I4–I5 and I5–I6 distances of 2.9493(5) and 2.8918(5) Å, respectively, and I4–I5–I6 angle is 172.954(17)°. The crystal packing of title compound is stabilized via multiple intermolecular contacts between $[\text{C}_{12}\text{H}_9\text{I}_3\text{NO}]^+$ cation and iodine atoms that can be attributed to halogen bonding [9]. The inversely oriented organic cations are linked through I1...I2' (3.8718(5) Å; ' = $-x+1, -y+1, -z$) intermolecular contacts forming dimeric structure associations. Additionally, these dimers are extended along *a*-axis direction through I2...I4' (3.4947(5) Å; ' = $-x+1, -y+1, -z$) contacts. Finally, the overall packing is completed by intermolecular contacts C10–H10B...I4' (3.1625(4) Å; ' = $-x+1, -y+1, -z$) and C10–H10B...I6' (3.1376(4) Å; ' = $x-1/2, -y+1/2+1, +z-1/2$).

* Correspondence author (e-mail: sgg@uniovi.es)

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(4)	4e	0.5307	0.5802	0.0045	0.019
H(7)	4e	1.0081	0.6186	0.1353	0.023
H(8)	4e	1.1518	0.7264	0.1567	0.026
H(9)	4e	1.0636	0.8470	0.1047	0.024
H(10A)	4e	0.6661	0.9312	−0.0950	0.023

Table 2. continued.

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(10B)	4e	0.7506	0.8581	−0.1344	0.023
H(11)	4e	0.8952	0.9226	−0.0027	0.019
H(12A)	4e	0.8222	0.9250	0.1835	0.023
H(12B)	4e	0.6653	0.9037	0.1386	0.023

Table 3. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₁₂	<i>U</i> ₁₃	<i>U</i> ₂₃
C(1)	4e	0.7782(5)	0.7490(3)	0.0389(4)	0.015(3)	0.012(3)	0.019(3)	−0.001(2)	0.005(2)	−0.001(2)
C(2)	4e	0.6375(6)	0.7601(3)	−0.0089(4)	0.019(3)	0.014(3)	0.014(3)	0.005(2)	0.008(2)	0.002(2)
C(3)	4e	0.5465(5)	0.6970(3)	−0.0204(4)	0.011(2)	0.019(3)	0.021(3)	0.003(2)	0.004(2)	−0.001(2)
C(4)	4e	0.5938(5)	0.6221(3)	0.0120(4)	0.019(3)	0.014(3)	0.014(3)	−0.002(2)	0.004(2)	0.000(2)
C(5)	4e	0.7311(6)	0.6105(3)	0.0544(4)	0.022(3)	0.015(3)	0.020(3)	0.004(2)	0.006(2)	0.002(2)
C(6)	4e	0.8286(5)	0.6738(3)	0.0711(4)	0.015(3)	0.012(3)	0.018(3)	0.002(2)	0.004(2)	0.004(2)
C(7)	4e	0.9718(6)	0.6672(3)	0.1148(4)	0.020(3)	0.018(3)	0.020(3)	0.002(2)	0.003(2)	0.001(2)
C(8)	4e	1.0577(6)	0.7312(3)	0.1272(4)	0.018(3)	0.026(3)	0.021(3)	0.001(2)	−0.002(2)	−0.002(2)
C(9)	4e	1.0045(6)	0.8035(3)	0.0960(4)	0.019(3)	0.020(3)	0.021(3)	−0.004(2)	0.002(2)	−0.007(2)
C(10)	4e	0.7049(6)	0.8809(3)	−0.0726(4)	0.024(3)	0.015(3)	0.017(3)	−0.002(2)	−0.001(2)	0.002(2)
C(11)	4e	0.8149(6)	0.8912(3)	0.0215(4)	0.021(3)	0.010(3)	0.018(3)	0.000(2)	0.000(2)	0.001(2)
C(12)	4e	0.7537(6)	0.9298(3)	0.1210(4)	0.021(3)	0.015(3)	0.022(3)	0.003(2)	−0.001(2)	0.004(2)
I(1)	4e	0.79737(4)	0.49600(2)	0.08560(3)	0.0272(2)	0.0139(2)	0.0259(2)	0.0030(2)	0.0041(2)	0.0036(2)
I(2)	4e	0.34078(4)	0.71640(2)	−0.09376(3)	0.0157(2)	0.0150(2)	0.0277(2)	−0.0010(1)	−0.0001(2)	0.0003(2)
I(3)	4e	0.71013(4)	1.05059(2)	0.08915(3)	0.0381(2)	0.0146(2)	0.0303(2)	0.0053(2)	−0.0011(2)	0.0005(2)
I(4)	4e	0.99358(4)	0.23970(2)	0.22474(3)	0.0190(2)	0.0294(2)	0.0227(2)	−0.0009(2)	0.0014(2)	0.0022(2)
I(5)	4e	1.17709(4)	0.37990(2)	0.23333(3)	0.0179(2)	0.0247(2)	0.0173(2)	0.0049(2)	0.0007(2)	−0.0007(2)
I(6)	4e	1.38620(4)	0.50472(2)	0.23675(3)	0.0301(2)	0.0257(2)	0.0287(2)	−0.0033(2)	0.0068(2)	−0.0082(2)
N(1)	4e	0.8694(5)	0.8122(2)	0.0533(3)	0.018(2)	0.015(2)	0.018(2)	−0.004(2)	0.005(2)	−0.001(2)
O(1)	4e	0.5906(4)	0.8314(2)	−0.0407(3)	0.016(2)	0.011(2)	0.026(2)	−0.001(2)	−0.003(2)	0.002(2)

Acknowledgments. We thank for financial support from the Spanish Ministerio de Economía y Competitividad under projects MAT2010-15094 and FEDER. A.D. also thanks to Spanish Ministerio de Educación, Cultura y Deporte for the FPU grant AP2008-03942. This work was supported by the Russian Foundation for Basic Research, grant 14-03-00961a. We also like to thank Joaquin Garcia Alvarez for the help in the realization of the synthesis.

References

- Kim, D. G.: Halocyclization of 8-allylthioquinoline. *Chem. Heterocycl. Compd.* **33** (1997) 1133-1135.
- Paira, R.; Paira, P.; Maity, A.; Mondal, S.; Hazra, A.; Sahu, K. B.; Naskar, S.; Saha, P.; Banerjee, M.; Mondal, N. B.: Amberlite IRA 402(OH): an efficient mediator for the exclusive synthesis of fused tricyclic oxaza quinolinium salts. *Tetrahedron Lett.* **51** (2010) 3200-3204.
- Kalita, D.; Baruah, J. B.: Rearrangement of 2-Bromo-*N*-quinoline-8-yl-acetamide Leading to New Heterocycle. *J. Heterocyclic Chem.* **47** (2010) 459-462.
- Mammadova, G. Z.; Matsulevich, Z. V.; Borisova, G. N.; Borisov, F. V.; Khrestalev, V. N.: (7a*R**,12b*S**)-8,12b-Dihydro-7a*H*-indeno[1',2':5,6]-[1,4]selenazino [2,3,4-*ij*]-quinolin-13-ium hydrogen sulfate. *Acta Crystallogr.* **E67** (2011) o3286-o3287.
- Slepukhin, P. A.; Batalov, V. I.; Kim, D. G.; Charushin, V. N.: A single crystal X-ray study of the products of halogen mercury cyclization of 8-allylthioquinoline. *J. Struct. Chem.* **52** (2012) 145-150.
- Svensson, P. H.; Kloo, L.: Synthesis, Structure, and Bonding in Polyiodide and Metal Iodide-Iodine Systems. *Chem. Rev.* **103** (2003) 1649-1684.
- Metrangolo, P.; Resnati, G.; Pilati, T.; Liantonio, R.; Meyer, F.: Engineering functional materials by halogen bonding. *J. Polym. Sci. Part A: Polym. Chem.* **45** (2007) 1-15.
- van Megen, M.; Reiss, G. J.: I₆^{2−} Anion Composed of Two Asymmetric Triiodide Moieties: A Competition between Halogen and Hydrogen Bond. *Inorganics* **1** (2013) 3-13 and references cited therein.
- Desiraju, G. R.; Ho, P. S.; Kloo, L.; Legon, A. C.; Marquardt, R.; Metrangolo, P.; Politzer, P. A.; Resnati, G.; Rissanen, K.: Definition of the halogen bond. *Pure Appl. Chem.* **85** (2013) 1711-1713.
- Agilent Technologies: CrysAlis PRO Software system, version 1.171.35.15, Agilent Technologies UK Ltd, Oxford, UK 2011.
- Altomare, A.; Casciarano, G.; Giacovazzo, C.; Guagliardi, A.; Burla, M. C.; Polidori, G.; Camalli, M.: SIR92 - a program for automatic solution of crystal structures by direct methods. *J. Appl. Cryst.* **27** (1994) 435-436.
- Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr.* **A64** (2008) 112-122.
- Brandenburg, K.: DIAMOND. Visual Crystal Structure Information System. Version 3.2i. Crystal Impact, Bonn, Germany 2012.
- Farrugia, L. J.: WinGX suite for small-molecule single-crystal crystallography. *J. Appl. Cryst.* **32** (1999) 837-838.
- Allen, F. H.; Johnson, O.; Shields, G. P.; Smith, B. R.; Towler, M.: CIF applications. XV. enCIFer: a program for viewing, editing and visualizing CIFs. *J. Appl. Cryst.* **37** (2004) 335-338.